

FOX INSPECTIONS 8616 DAFFODIL ST HOUSTON, TX 77063

Certificate of Mold Analysis

Prepared for:	FOX INSPECTIONS
Phone Number:	(713) 723-3330
Fax Number:	
Project Name:	Erick & Mara Calderon
Test Location:	90689 - 210604AD
-2135 Quenby Street,	

Report Number: Received Date: Report Date: Houston, TX 77005 1434424 June 7, 2021 June 7, 2021

Ina

Diana Sauri, Laboratory Director or other approved signatory

Currently there are no Federal regulations for evaluating potential health effects of fungal contamination and remediation. This information is subject to change as more information regarding fungal contaminants available visit becomes For more information http://www.epa.gov/mold or www.nyc.gov/html/doh/html/epi/mold.shtml. This document was designed to follow currently known industry guidelines for the interpretation of microbial sampling, analysis, and remediation. Since interpretation of mold analysis reports is a scientific work in progress, it may as such be changed at any time without notice. The client is solely responsible for the use or interpretation. PRO-LAB/SSPTM Inc. makes no express or implied warranties as to health of a property from only the samples sent to their laboratory for analysis. The Client is hereby notified that due to the subjective nature of fungal analysis and the mold growth process, laboratory samples can and do change over time relative to the originally sampled material. PRO-LAB/SSPTM Inc. reserves the right to properly dispose of all samples after the testing of such samples are sufficiently completed or after a 7 day period, whichever is greater.



For more information please contact PRO-LAB at (954) 384-4446 or email info@prolabinc.com



1675 North Commerce Parkway, Weston, FL 33326 (954) 384-4446

Prepared for : FOX INSPECTIONS

Test Address : Erick & Mara Calderon 90689 - 210604AD

-2135 Quenby Street,

Houston, T	X 77005

ANALYSIS METHOD	6110 Air Direct Examination		6110 Air Direct Examination		6110 Air Direct Examination			6110 Air Direct Examination				
LOCATION	OUTSIDE		LIVING ROOM		OFFICE (BY SKYLIGHT)		THIRD FLOOR STAIRS					
COC / LINE #	1434424 - 1		1434424 - 2		1434424 - 3		1434424 - 4					
SAMPLE TYPE & VOLUME	AIR-O-CELL - 150.00L		AIR-O-CELL - 150.00L		AIR-O-CELL - 150.00L		AIR-O-CELL - 150.00L					
SERIAL NUMBER	31993036		31993054			31993078			31993052			
COLLECTION DATE	Jun 4, 2021		Jun 4, 2021			Jun 4, 2021			Jun 4, 2021			
ANALYSIS DATE	Jun 7, 2021		Jun 7, 2021			Jun 7, 2021			Jun 7, 2021			
CONCLUSION	CONTROL		NOT ELEVATED			NOT ELEVATED			NOT ELEVATED			
IDENTIFICATION	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total
Cladosporium	28	190	29	12	80	23	4	27	34	36	240	41
Other Ascospores	40	270	42									
Other Basidiospores	12	80	12									
Penicillium/Aspergillus	16	110	17	40	270	77	8	53	66	44	290	50
Spegazzinia										8	53	9
TOTAL SPORES	96	650	100	52	350	100	12	80	100	88	583	100
MINIMUM DETECTION LIMIT*	4	27		4	27		4	27		4	27	
BACKGROUND DEBRIS	Moderate		Light		Light			Moderate				
Cellulose Fiber	16	110					4	27		12	80	
OBSERVATIONS & COMMENTS	Non-biological debris present.									Non-biological debris present.		

Background debris qualitatively estimates the amount of particles that are not pollen or spores and directly affects the accuracy of the spore counts. The categories of Light, Moderate, Heavy and Too Heavy for Accurate Count, are used to indicate the amount of deposited debris. Light (None to up to 25% obstruction); Medium (26% to up to 75% obstruction); Heavy (76% to up to 90% obstruction); Too Heavy (Greater than 90% obstruction). Increasing amounts of debris will obscure small spores and can prevent spores from impacting onto the slide. The actual number of spores present in the sample is likely higher than reported if the debris estimate is 'Heavy' or 'Too Heavy for Accurate Count'. All calculations are rounded to two significant figures and therefore, the total percentage of spore numbers may not equal 100%. The effect of the results relate only to the items tested. The methods used in this analysis have been validated and is fit for the intended use. R "version" indicated after the lab ID# indicates a sample with amended data.

Minimum Detection Limit. Based on the volume of air sampled, this is the lowest number of spores that can be detected and is an estimate of the lowest concentration of spores that can be read in the sample. NA = Not Applicable

Spores that were observed from the samples submitted are listed on this report. If a spore is not listed on this report it was not observed in the samples submitted.

Interpretation Guidelines: A determination is added to the report to help users interpret the mold analysis results. A mold report is only one aspect of an indoor air quality investigation. The most important aspect of mold growth in a living space is the availability of water. Without a source of water, mold generally will not become a problem in buildings. These determinations are in no way meant to imply any health outcomes or financial decisions based solely on this report. For questions relating to medical conditions you should consult an occupational or environmental health physician or professional. **CONTROL** is a baseline sample showing what the spore count and diversity is at the time of sampling. The control sample(s) is usually collected outside of the structure being tested and used to determine if this sample(s) is similar in diversity and abundance to the inside sample(s). ELEVATED means that the amount and/or diversity of spores, as compared to the control sample(s), and other samples in our database, are higher than expected. This can indicate that fungi have grown because of

a water leak or water intrusion. Fungi that are considered to be indicators of water damage include, but are not limited to: Chaetomium, Fusarium, Memnoniella, Stachybotrys, Scopulariopsis, Ulocladium. NOT ELEVATED means that the amount and/or the diversity of spores, as compared to the control sample and other samples in our database, are lower than expected and may indicate no problematic fungal growth. UNUSUAL means that the presence of current or former growth was observed in the analyzed sample. An abundance of spores are present, and/or growth structures including hyphae and/or fruiting bodies are Present and associated with one or more of the types of mold/lungi identified in the analyzed sample. NORMAL means that no presence of current or former growth was observed in the analyzed sample.



1675 North Commerce Parkway, Weston, FL 33326 (954) 384-4446

Prepared for : FOX INSPECTIONS

Test Address : Erick & Mara Calderon 90689 - 210604AD

-2135 Quenby Street,

Houston, TX 77005

ANALYSIS METHOD	6110 Air Direct Examination		INTENTIONALLY BLANK			INTENTIONALLY BLANK			INTENTIONALLY BLANK			
LOCATION	MASTER BEDROOM											
COC / LINE #	1434424 - 5											
SAMPLE TYPE & VOLUME	AIR-0	D-CELL - 15	0.00L									
SERIAL NUMBER		31993055										
COLLECTION DATE	Jun 4, 2021											
ANALYSIS DATE	Jun 7, 2021											
CONCLUSION	NOT ELEVATED											
IDENTIFICATION	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total	Raw Count	Spores per m ³	Percent of Total
Cladosporium												
Other Ascospores												
Other Basidiospores												
Penicillium/Aspergillus	36	240	100									
Spegazzinia												
TOTAL SPORES	36	240	100									
MINIMUM DETECTION LIMIT*	4	27										
BACKGROUND DEBRIS	Light											
Cellulose Fiber	16	110										
OBSERVATIONS & COMMENTS												


Background debris qualitatively estimates the amount of particles that are not pollen or spores and directly affects the accuracy of the spore counts. The categories of Light, Moderate, Heavy and Too Heavy for Accurate Count, are used to indicate the amount of deposited debris. Light (None to up to 25% obstruction); Medium (26% to up to 75% obstruction); Heavy (76% to up to 90% obstruction); Too Heavy (Greater than 90% obstruction). Increasing amounts of debris will obscure small spores and can prevent spores from impacting onto the slide. The actual number of spores present in the sample is likely higher than reported if the debris estimate is 'Heavy' or 'Too Heavy for Accurate Count'. All calculations are rounded to two significant figures and therefore, the total percentage of spore numbers may not equal 100%. The effect of the results relate only to the items tested. The methods used in this analysis have been validated and is fit for the intended use. R "version" indicated after the lab ID# indicates a sample with amended data.


Minimum Detection Limit. Based on the volume of air sampled, this is the lowest number of spores that can be detected and is an estimate of the lowest concentration of spores that can be read in the sample. NA = Not Applicable


Spores that were observed from the samples submitted are listed on this report. If a spore is not listed on this report it was not observed in the samples submitted.


Interpretation Guidelines: A determination is added to the report to help users interpret the mold analysis results. A mold report is only one aspect of an indoor air quality investigation. The most important aspect of mold growth in a living space is the availability of water. Without a source of water, mold generally will not become a problem in buildings. These determinations are in no way meant to imply any health outcomes or financial decisions based solely on this report. For questions relating to medical conditions you should consult an occupational or environmental health physician or professional. **CONTROL** is a baseline sample showing what the spore count and diversity is at the time of sampling. The control sample(s) is usually collected outside of the structure being tested and used to determine if this sample(s) is similar in diversity and abundance to the inside sample(s). ELEVATED means that the amount and/or diversity of spores, as compared to the control sample(s), and other samples in our database, are higher than expected. This can indicate that fungi have grown because of

a water leak or water intrusion. Fungi that are considered to be indicators of water damage include, but are not limited to: Chaetomium, Fusarium, Memnoniella, Stachybotrys, Scopulariopsis, Ulocladium. NOT ELEVATED means that the amount and/or the diversity of spores, as compared to the control sample and other samples in our database, are lower than expected and may indicate no problematic fungal growth. UNUSUAL means that the presence of current or former growth was observed in the analyzed sample. An abundance of spores are present, and/or growth structures including hyphae and/or fruiting bodies are Present and associated with one or more of the types of mold/lungi identified in the analyzed sample. NORMAL means that no presence of current or former growth was observed in the analyzed sample.











1675 North Commerce Parkway, Weston, FL 33326 (954) 384-4446

Identification	Outdoor Habitat	Indoor Habitat	Possible Allergic Potential Not an opinion or interpretation	Comments
Cladosporium	The most common spore type reported in the air worldwide. Found on dead and dying plant litter, and soil.	Commonly found on wood and wallboard. Commonly grows on window sills, textiles and foods.	Type I (hay fever and asthma), Type III (hypersensitivity pneumonitis) allergies.	A very common and important allergen source both outdoors and indoors.
Ascospores	Common everywhere. Constitutes a large part of the airspora outside. Can reach very high numbers in the air outside during the spring and summer. Can increase in numbers during and after rainfalls.	Very few of this group grow inside. The notable exception is Chaetomium, Ascotricha and Peziza.	Little known for most of this group of fungi. Dependent on the type (see Chaetomium and Ascotricha).	
Basidiospores	Commonly found everywhere, especially in the late summer and fall. These spores are from Mushrooms.	Mushrooms are not normally found growing indoors, but can grow on wet lumber, especially in crawlspaces. Sometimes mushrooms can be seen growing in flower pots indoors.	Some allergenicity reported. Type I (hay fever, asthma) and Type III (hypersensitivity pneumonitis).	Among the group of Mushrooms (Basidiomycetes) are dry rot fungi Serpula and Poria that are particularly destructive to buildings.
Penicillium/Aspergillus	Common everywhere. Normally found in the air in small amounts in outdoor air. Grows on nearly everything.	Wetted wallboard, wood, food, leather, etc. Able to grow on many substrates indoors.	Type I (hay fever and asthma) allergies and Type III (hypersensitivity pneumonitis) allergies.	This is a combination group of Penicillium and Aspergillus and is used when only the spores are seen. The spores are so similar that they cannot be reliably separated into their respective genera.
Spegazzinia	Not commonly observed, but widely distributed.	Not known to grow indoors.	None known.	Frequently seen especially in southern United States.